

## HERBICIDE

# TRIAD SELECT™

SELECTIVE, BROAD-SPECTRUM WEED CONTROL.



**Triad Select™** utilizes a combination of three active ingredients to deliver control of a wide variety of hard-to-kill broadleaf weeds such as dandelion, clover, and plantain.

**Triad Select** has been demonstrated through university trials to be an effective solution that can be applied to both cool- and warm-season grasses, and is labeled for use on turf, sod farms, rights-of-ways, and other non-crop industrial areas.

## PRODUCT FEATURES

- Combined power of three selective herbicides.
- Low-odor formulation with flexible application rates.
- For use on both cool- and warm-season grasses.
- May be applied to woody brush in non-cropland areas.
- Can be tank-mixed with liquid fertilizers or liquid iron materials (Refer to label for fertilizer tank-mix compatibility.)
- University-tested.

## LABELED USES

- Golf courses
- Cemeteries
- Parks & sports fields
- Turfgrass, lawns & grass
- Turf grown for seed & sod
- Fence & hedgerows
- Roadsides & ditches
- Rights-of-way
- Utility power lines
- Railroads, airports & industrial sites

See label for complete application guidelines and restrictions.

### ACTIVE INGREDIENT

30.89% 2,4-D  
8.23% MCPA  
2.77% dicamba

### HRAC GROUP

2,4-D - Group 4  
MCPA - Group 4  
Dicamba - Group 4

### FORMULATION

Liquid concentrate

### CHEMICAL FAMILY

Phenoxy-carboxylic acids  
Benzoic acids

### COMPARE TO

Proprietary formulation

### MODE OF ACTION

Auxin mimic

### RESTRICTED USE

No

### SIGNAL WORD

Warning

### PACKAGING

12 x 1 qt, 4 x 1 gal, 2 x 2.5 gal,  
5 x 30 gal drum, 55 gal drum,  
265 gal tote

### EPA REG NUMBER

89442-22

# TRIAD SELECT™

SELECTIVE, BROAD-SPECTRUM WEED CONTROL.

## APPLICATION INFORMATION

### Non-turf application rate

**Roadsides (including aprons and guardrails), rights-of-way, and other similar non-crop areas:** For control of broadleaf weeds, mix at a rate of 1.5 to 3.39 quarts of product per acre. This may require 50 to 300 gallons of water per acre (43,560 sq ft). For woody plants, apply to both stems and foliage of completely matured plants until they start to go dormant.

**Woody Brush:** Add 1 to 1.5 gallons of this product per acre to wet all parts of the brush foliage, stem, and bark. This may require up to 200 to 600 gallons of water per acre (43,560 sq ft).

### Bahiagrass, bluegrass, common bermudagrass, fescue, ryegrass, zoysia grass

Use Rate Fluid Ounces	Spray Volume Gallons	Use Rate Fluid Ounces	Spray Volume Gallons
Per 1,000 sq ft		Per Acre	
1.1 to 1.5	0.5 to 5	3 to 4	5 to 220

### Dormant St. Augustine

(Common, Raleigh, and Seville varieties grown in Texas, Louisiana, and Mississippi only)

Use Rate Fluid Ounces	Spray Volume Gallons	Use Rate Fluid Ounces	Spray Volume Gallons
Per 1,000 sq ft		Per Acre	
0.75 to 0.90 maximum	1 to 5	2 to 2.5 maximum	40 to 200

### Bentgrass

(Putting and bowling greens)

Use Rate Fluid Ounces	Spray Volume Gallons	Use Rate Fluid Ounces	Spray Volume Gallons
Per 1,000 sq ft		Per Acre	
0.67 maximum	3.3	1.8 maximum	145




◀ See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. Albaugh and the A logo are registered trademarks and Triad Select is a trademark of Albaugh, LLC. Trimec is a registered trademark of PBI-Gordon Corporation. Quali-pro is a registered trademark of Adama. Always read and follow label restrictions, precautions, and directions for use. This product may not be registered for sale or use in all states or counties. Please check with your state agency responsible for pesticide registration to ensure registration status.


### Virginia Tech - Cool Weather Trial

% Percent control @ 16 DAT  
Applied to Tall fescue, Kentucky bluegrass, and Perennial ryegrass  
Untreated = 0% Control



### Clemson University - Warm Weather Trial

% Percent control @ 35 DAT  
Applied to: Bermuda, Tall fescue • Untreated = 0% Control



## LABELED WEEDS

- |             |                  |                                   |
|-------------|------------------|-----------------------------------|
| Bedstraw    | Knotweed         | Ragweed                           |
| Black Medic | Lambsquarters    | Sheep Sorrel                      |
| Buckhorn    | <i>Lespedeza</i> | Shepherds purse                   |
| Chicory     | Mallow           | Speedwell                         |
| Chickweed   | Morning glory    | Spurge                            |
| Clover      | Peppergrass      | Wild Carrot                       |
| Dandelion   | Pigweed          | Wild Garlic                       |
| Dock        | Plantain         | Wild Lettuce                      |
| Ground Ivy  | Poison Ivy       | Wild Onion                        |
| Heal-All    | Poison Oak       | Yarrow & other<br>broadleaf weeds |
| Henbit      | Purslane         |                                   |